

Risk	Indicators For Increased Risk	Contributing Factors	Possible Interventions
High Acute Risk	<ul style="list-style-type: none"> * Persistent suicidal ideation or thought *Strong intention to act or plan, OR *Not able to control Impulse 	<ul style="list-style-type: none"> *Acute state of psychiatric disorder or acute psychiatric symptoms *Acute precipitating event(s) *Inadequate protective factors 	<p>Direct soldier to Emergency room</p> <p>Admission generally indicated, unless significant change reduces risk.</p>
Intermediate Acute Risk	<ul style="list-style-type: none"> *Current Suicidal ideation or thoughts *No intent to act *Able to control impulse *No recent attempt or preparatory behaviors or rehearsal of act 	<ul style="list-style-type: none"> *Existence of warning signs or risk factors; <p>AND</p> <ul style="list-style-type: none"> *Limited protective factors 	<p>Emergency room may be necessary depending on risk factors. Develop/Safety plan. Give emergency/crisis numbers, and connect soldier with PHP immediately</p>
Low Acute Risk	<ul style="list-style-type: none"> *Recent suicidal ideation or thoughts *No intention to act on plan, able to control impulse *No planning or rehearsing a suicide act *No previous attempt 	<ul style="list-style-type: none"> *Existence of protective factors *Limited risk factors 	<p>Make a referral to PHP, and request update on community or military services/resources to aid in symptom reduction. Give emergency/crisis numbers.</p>
Undetermined Risk	<p>Due to difficulty in determining the level of risk, Unit has concerns about soldier despite denial of ideation or intent. The soldier should be immediately referred for assistance by PHP.</p>		